

Dynamical study of the possible molecular state $X(3872)$ with the s -channel one gluon exchange interaction

Bao-Kai Wang and Wei-Zhen Deng*

*Department of Physics and State Key Laboratory of Nuclear Physics and Technology,
Peking University, Beijing 100871, China*

Xiao-Lin Chen

Department of Physics, Peking University, Beijing 100871, China

Abstract

The recently observed $X(3872)$ resonance, which is difficult to be assigned a conventional $c\bar{c}$ charmonium state in the quark model, may be interpreted as a molecular state. Such a molecular state is a hidden flavor four quark state because of its charmonium-like quantum numbers. The s -channel one gluon exchange is an interaction which only acts in the hidden flavor multi-quark system. In this paper, we will study the $X(3872)$ and other similar hidden flavor molecular states in a quark model by taking into account of the s -channel one gluon exchange interaction.

*Electronic address: dwz@th.phy.pku.edu.cn

I. INTRODUCTION

Recently, many new resonances are discovered experimentally. Many of them have the proper quantum numbers of the $q\bar{q}$ meson states. However, their mass values do not fit the conventional $q\bar{q}$ states in the quark model. Among them, $X(3872)$ was first observed in the $J/\psi\pi^+\pi^-$ channel by Belle collaboration in 2003 [1], and has been confirmed by CDF [2], D0 [3] and Babar collaborations [4]. Its quantum numbers are probably $J^{PC} = 1^{++}$. The corresponding charmonium candidates in the quark model are $2^3P_1(3990)$ and $3^3P_1(4290)$ which are $50 \sim 200$ MeV above $M_X = 3872$ MeV.

Many people suggested that $X(3872)$ is mainly a $D\bar{D}^*$ molecular state [5, 6, 7, 8, 9]. However, to bind the quarks and anti-quarks together in such a four quark state or other multi-quark states, we need to introduce new interaction into the quark model. Swanson proposed that the $X(3872)$ is mainly a $D\bar{D}^*$ molecule bound by the meson-meson interaction derived from the one pion exchange and the quark exchange [7]. In Wong's work [8], the meson-meson interaction is derived from a QED-type effective interaction in terms of effective charges for quarks and antiquarks. In refs. [10, 11, 12, 13, 14, 15], further investigations based on the molecule assumption were carried out.

Since the strength of the one pion exchange interaction seems not strong enough to bind the $D\bar{D}^*$ molecular state, other authors argued that $X(3872)$ may be a dominant $c\bar{c}$ charmonium with some admixture of $D\bar{D}^*$ [16, 17, 18]. In ref. [19], after taking into account of the sigma meson exchange potential, the interpretation of $X(3872)$ as a loosely bound molecular state was further disfavored.

We should notice that the color structure of a multi-quark state is much richer than that of a $q\bar{q}$ conventional meson state. Unlike the conventional mesons or baryons, the $q\bar{q}$ and qq pairs in a multi-quark state can be in the color 8_c and 6_c representations respectively. Some color interactions which have no effects in the $q\bar{q}$ or q^3 colorless system, may have significant contribution in a multi-quark system. So the complete interactions in the quark model can be quite different after we take into account of these multi-quark states.

The s -channel one gluon exchange interaction is an interaction between quark and anti-quark of the same flavor which annihilate into a virtual gluon. It has no effect on the conventional $q\bar{q}$ mesons but acts in the hidden flavor multi-quark system like the charmonium-like molecular states. In this work, we will investigate the hidden flavor molecular states by con-

sidering the s -channel gluon exchange interaction in the quark model. In the next section, we will model the potential of the s -channel one gluon exchange interaction starting from the non-relativistic reduction. Then the J^{PC} quantum numbers of the molecular states are selected by an analysis of the spin dependence of the interaction strength. In sec. III, we will carry out the numerical calculation of $X(3872)$ and some other charmonium-like states as the molecular states. Also we will make prediction about the similar bottomium-like molecular states. Finally, we will give a brief summary.

II. THE POTENTIAL OF S CHANNEL ONE GLUON EXCHANGE INTERACTION

In our work, we will use the Bhaduri quark model which is a rather simple non-relativistic quark potential model. In the Bhaduri model, the hamiltonian can be written as[20]

$$H = \sum_i (m_i + \frac{\mathbf{P}_i^2}{2m_i}) - \frac{3}{4} \sum_{i < j} (\mathbf{F}_i \cdot \mathbf{F}_j V_{ij}^C + \mathbf{F}_i \cdot \mathbf{F}_j \mathbf{S}_i \cdot \mathbf{S}_j V_{ij}^{SS}). \quad (1)$$

m_i are constituent quark masses and $\mathbf{F}_i^c = \frac{1}{2} \boldsymbol{\lambda}_i^c$ ($c = 1, \dots, 8$) are the well-known $SU_c(3)$ Gell-Mann matrices. Apart from a constant, here the central potential is the usual one gluon exchange coulomb potential plus the linear confinement:

$$V_{ij}^C = -\frac{\kappa}{r_{ij}} + \frac{r_{ij}}{a_0^2} - M_0. \quad (2)$$

$r_{ij} = |\mathbf{r}_i - \mathbf{r}_j|$ is the distance between quark i and j . The color-magnetic interaction reads

$$V_{ij}^{SS} = \frac{4\kappa}{m_i m_j} \frac{1}{r_0^2 r_{ij}} e^{-r_{ij}/r_0}, \quad (3)$$


where the δ -interaction has been smeared smoothly with the prescription

$$\delta^3(\mathbf{r}) \rightarrow \frac{1}{4\pi r_0^2 r_{ij}} e^{-r_{ij}/r_0}. \quad (4)$$

The model parameter values are

$$\begin{aligned} \kappa &= 102.67 \text{MeVfm}, & a_0 &= 0.0326(\text{MeV}^{-1}\text{fm})^{\frac{1}{2}}, \\ M_0 &= 913.5 \text{MeV}, & r_0 &= 0.4545 \text{fm} \\ m_u &= m_d = 337 \text{MeV}, & m_s &= 600 \text{MeV}, \\ m_c &= 1870 \text{MeV}, & m_b &= 5259 \text{MeV}. \end{aligned}$$

FIG. 1: The s -channel one gluon exchange



The s -channel one gluon exchange interaction (SOGEx) takes place when a $q\bar{q}$ pair annihilates into a virtual gluon (Fig. 1). The non-relativistic reduction of the potential is:

$$V^{\text{SOGEx}}(\mathbf{r}_{ij}) = -\frac{1}{2} \left(\frac{4}{3} + \mathbf{F}_q \cdot \mathbf{F}_{\bar{q}} \right) \left(1 + \frac{4}{3} \mathbf{S}_q \cdot \mathbf{S}_{\bar{q}} \right) G(4m^2) \delta^3(\mathbf{r}_{ij}) \delta(f_i, f_j), \quad (5)$$

where $G(4m^2)$ is the one gluon exchange amplitude. The last $\delta(f_i, f_j)$ represents that the quark and the anti-quark must have the same flavor. Clearly, the two factors in the brackets mean that this interaction only occurs when the $q\bar{q}$ pair is in the color octet and with spin $S = 1$. In our model calculation, the δ -interaction should be smeared smoothly with the same prescription (4) used in the color-magnetic interaction in the same quark model. However, the QCD amplitude $G(4m^2)$ is taken value in the timelike region, where the QCD behavior is still not clear at present time. In the spacelike region, this amplitude is well known from the gluon propagator in perturbative QCD and it reads

$$G_{\text{pert}}^t(q^2) = \frac{4\pi\alpha_s}{-q^2} \quad \text{for } q^2 < 0. \quad (6)$$

However, in our study of the hidden flavor molecular states, in order to provide an attractive interaction to favor the formation of the bound states, we need that

$$G(4m^2) > 0,$$

which means the above formula from perturbative QCD should not be directly used in the timelike region. This change of sign is first suggested in ref. [21] in the study of $\pi\pi$ and $K\pi$ scattering. Following ref. [21], we assume that

$$G(4m^2) = -f G_{\text{pert}}^t(4m^2) = f \frac{\pi\alpha_s^2}{m^2}, \quad (7)$$

where f is an introduced strength factor. In our model, after the δ -function being smeared smoothly, the SOGE potential turns to be:

$$V_{ij}^{\text{SOGE}} = -\frac{f}{4} \left(1 + \frac{3}{4} \mathbf{F}_q \cdot \mathbf{F}_{\bar{q}} \right) \left(\frac{3}{4} + \mathbf{S}_q \cdot \mathbf{S}_{\bar{q}} \right) \frac{4\kappa}{m_i^2 r_0^2 r_{ij}} e^{-r_{ij}/r_0} \delta(f_i, f_j). \quad (8)$$

First we observe that the potential V^{SOGE} is proportional to m_i^{-2} , so the interaction mainly takes place between the light $q\bar{q}$ pairs of $q = u, d, s$ in any multi-quark system.

Next, we will analyse the spin dependence of V^{SOGE} in each J^{PC} channel of molecular state. For the molecular states we concern, let us assume that their flavor structure is $Q\bar{q}q\bar{Q}$, where $Q = c, b$ and $q = u, d, s$. Since we can neglect the V^{SOGE} interaction between $Q\bar{Q}$, the interaction acts only when the $q\bar{q}$ pair is in color octet and $S = 1$ as we have mentioned before. More specifically, the interaction will be switched on if the spin coupling of the four quarks is

$$[(Q\bar{Q})_{J_1}(q\bar{q})_{J_2=1}]_J.$$

The color structure of $Q\bar{Q}$ and $q\bar{q}$ should be octet obviously. However, the color (re)coupling is irrelevant to our analysis here and will not be explicitly presented. Furthermore, we will assume no spatial excitation of any quark. Hence all quarks have zero orbital angular momentum. Then the quantum numbers J^{PC} can be determined easily for $J_1 = 0, 1$. We have following four V^{SOGE} interaction channels:

$$J_1 = 0 : 1^{+-}; \quad J_1 = 1 : 0^{++}, 1^{++}, 2^{++}.$$

Finally, we can recover the molecular states by the angular momentum recoupling. We obtain

- 0^{++}

$$[(Q\bar{Q})_1(q\bar{q})_1]_0 = -\frac{\sqrt{3}}{2}(Q\bar{q})_0(q\bar{Q})_0 - \frac{1}{2}[(Q\bar{q})_1(q\bar{Q})_1]_0,$$

- 1^{++}

$$[(Q\bar{Q})_1(q\bar{q})_1]_1 = -\frac{1}{\sqrt{2}}(Q\bar{q})_1(q\bar{Q})_0 + \frac{1}{\sqrt{2}}(Q\bar{q})_0(q\bar{Q})_1,$$

- 1^{+-}

$$(Q\bar{Q})_0(q\bar{q})_1 = \frac{1}{2}(Q\bar{q})_1(q\bar{Q})_0 + \frac{1}{2}(Q\bar{q})_0(q\bar{Q})_1 + \frac{1}{\sqrt{2}}[(Q\bar{q})_1(q\bar{Q})_1]_1,$$

TABLE I: The list of spin factor of V^{SOGE} interaction strength in a molecular state J^{PC} made of two meson states $J_1^{P_1}$ and $J_2^{P_2}$. *c.c.* = charge conjugation.

$J_1^{P_1} J_2^{P_2}$	J^{PC}	Spin factor
$0^- 0^-$	0^{++}	$\frac{3}{4}$
$1^- 1^-$	0^{++}	$\frac{1}{4}$
$1^- 0^- + c.c.$	1^{++}	1
$1^- 0^- - c.c.$	1^{+-}	$\frac{1}{2}$
$1^- 1^-$	1^{+-}	$\frac{1}{2}$
$1^- 1^-$	2^{++}	1

- 2^{++}

$$[(Q\bar{Q})_1(q\bar{q})_1]_2 = [(Q\bar{q})_1(q\bar{Q})_1]_2$$

Then the factor of interaction strength can be read from the coefficients, which we present in table I. We see that the V^{SOGE} interaction favors to form the molecular states with $J^{PC} = 1^{++}, 2^{++}$.

III. NUMERICAL CALCULATION

To calculate the molecular states, we use the Rayleigh-Ritz variation principle. The test wave function will be taken to be a series of Gaussian basis functions. The Gaussian basis functions are often utilized in variational calculations of atomic and molecular problems. Recently the method has been also used in the few body system in nuclear and particle physics [22, 23, 24].

In our case of the $Q\bar{q}q\bar{Q}$ molecular state, the test wave function of a molecular state between two clusters of $q\bar{q}$ meson states is a series

$$\psi_{1234}(r_{12}, r_{34}, r_{1234}) = \sum_i \alpha_{1234}^i \psi_{12}(r_{12}) \psi_{34}(r_{34}) \exp(-\beta_{1234}^i r_{1234}^2), \quad (9)$$

where $\mathbf{r}_1, \mathbf{r}_2, \mathbf{r}_3$ and $\mathbf{r}_4$ are the coordinates of $Q, \bar{q}, q$ and $\bar{Q}$, respectively. $\mathbf{r}_{ij} = \mathbf{r}_i - \mathbf{r}_j$.

r_{1234} is the distance between the two meson clusters

$$\mathbf{r}_{1234} = \frac{m_Q \mathbf{r}_1 + m_q \mathbf{r}_2}{m_Q + m_q} - \frac{m_q \mathbf{r}_3 + m_Q \mathbf{r}_4}{m_q + m_Q}. \quad (10)$$

$\psi_{ij}(r_{ij})$ is the meson wave function which is also taken to be a Gaussian function series

$$\psi_{ij}(r_{ij}) = \sum_k \alpha_{ij}^k \exp(-\beta_{ij}^k r_{ij}^2). \quad (11)$$

The wave function of a molecular state is determined by the variation principle in two steps. We first determine the wave function (11) of each meson cluster. Then the meson cluster functions ψ_{ij} are fixed in (9) to obtain the wave function of the molecular state and their masses.

To reduce the amount of computation, the parameters β^i and α^i in a Gaussian function series are determined in two steps by one-dimensional minimization. We first determined a average β value using a single Gaussian function. Then a set $\{\beta^i\}$ of $2N + 1$ elements is generated from scaling the β value up and down by a scale factor s [24]:

$$\beta^i = \beta s^{i-N} \quad (12)$$

where $i = 0, 1, \dots, 2N$. The coefficients α^i are determined by diagonalization the model Hamiltonian in the $2N + 1$ -dimensional space spanned by these $2N + 1$ different Gaussian functions. The final values of β^i and α^i and the mass of tetra quark states are determined by searching the scale factor s for a minimum of system energy.

In this way, we have calculated the possible 0^{++} (the combination of two 0^- mesons with the spin factor $\frac{3}{4}$), 1^{++} and 2^{++} molecular states in table I. We have calculated both the charmonium-like states $c\bar{q}q\bar{c}$ which will be compared to recent experimental results and the bottomium-like states $b\bar{q}q\bar{b}$ which can be investigated in further experiments.

The results of relevant heavy quark $Q\bar{q}$ mesons are listed in table II. We see that the mass values calculated from the Bhaduri quark model deviate at most 30 MeV away from the experimental data. The difference between the calculation from the variation method with Gaussian function series (cal. II) and the exact numerical calculation (cal. I) is less than 0.5 MeV. So the variation method is an impresssive good approximation for the numerical calculation of the $Q\bar{q}$ conventional mesons.

The results of the molecular states are given in table III and table IV. Here the molecular state is presented by its binding energy:

$$E_b = M_1 + M_2 - M_X, \quad (13)$$

TABLE II: Mass of $Q\bar{q}$ mesons. The experimental values are taken from PDG[25]. In calculation I, the mass is obtained by solving the Schrödinger equation. In calculation II, the mass is obtained by the variation method with Gaussian test functions using $N = 3$.

meson state	exp. (MeV)	cal. I (MeV)	cal. II (MeV)
$D^\pm$	1869.62	1885.56	1886.14
D^0	1864.84		
$D^*(2007)^0$	2006.97	2019.96	2020.07
$D^*(2010)^\pm$	2010.27		
$D_s^\pm$	1968.49	1995.78	1996.36
$D_s^{*\pm}$	2112.3	2101.2	2101.4
$B^\pm$	5279.15	5300.84	5301.18
B^0	5279.53		
B^*	5325.1	5350.3	5350.5
B_s^0	5366.3	5371.9	5372.4
B_s^*	5412.8	5413.3	5413.6

and the rms radius $\langle r^2 \rangle^{1/2}$, where M_1 , M_2 are masses of the two compound mesons, M_X is the mass of the molecular state.

We can see from tables III and IV that indeed the 1^{++} and 2^{++} molecular states are bound preferable since the V^{SOGE} are strong in these two channels. The binding of 1^{++} states are slightly deeper and tighter than that of 2^{++} 's. So from our model calculation, the 1^{++} molecular states should be easily observed in experiments.

First, let us look at the $D^{(*)}D^{(*)}$ sector. If the $X(3872)$ is a molecular state of D and D^* , the binding energy can be estimated from experimental data[25]

$$E_b = M_D + M_{D^*} - M_X = 0.3 \text{ MeV},$$

which is very small. Accounting the uncertainty of approximation in our model calculation, we think the reasonable range of f value should be around $-0.8 \sim -1.0$. If it is small $f \sim -0.8$, then it may be the only molecular states in the $D^{(*)}D^{(*)}$ sector. If $f \sim -1.0$, the 2^{++} state may also exists. From our numerical calculation, its mass should be about

$$M = M_{D^*} + M_{D^*} - E_b \approx 4008.5 \text{ MeV}.$$

TABLE III: Binding energy E_b (in MeV) of molecular states vs paramter f .

f	-0.8	-1.0	-1.5	-2.0	-3.0
$(DD)0^{++}$	—	—	11.8	28.1	71.7
$(DD^*)1^{++}$	1.8	6.2	24.2	48.7	109.3
$(D^*D^*)2^{++}$	—	5.4	21.4	43.5	98.1
$(D_sD_s)0^{++}$	—	—	—	—	5.3
$(D_sD_s^*)1^{++}$	—	—	—	—	15.4
$(D_s^*D_s^*)2^{++}$	—	—	—	—	14.0
$(BB)0^{++}$	8.8	15.3	35.2	58.5	110.9
$(BB^*)1^{++}$	16.8	26.8	55.8	88.5	160.4
$(B^*B^*)2^{++}$	16.0	25.5	53.4	84.7	153.6
$(B_sB_s)0^{++}$	—	—	3.4	9.3	25.7
$(B_sB_s^*)1^{++}$	—	1.9	8.9	18.9	44.4
$(B_s^*B_s^*)2^{++}$	—	1.8	8.5	18.2	42.9

TABLE IV: rms $\langle r^2 \rangle^{1/2}$ (in fm) of molecular states vs paramter f .

f	-0.8	-1.0	-1.5	-2.0	-3.0
$(DD)0^{++}$	—	—	1.33	1.00	0.76
$(DD^*)1^{++}$	2.67	1.68	1.07	0.86	0.69
$(D^*D^*)2^{++}$	—	1.79	1.12	0.91	0.72
$(D_sD_s)0^{++}$	—	—	—	—	1.63
$(D_sD_s^*)1^{++}$	—	—	—	—	1.12
$(D_s^*D_s^*)2^{++}$	—	—	—	—	1.17
$(BB)0^{++}$	1.11	0.94	0.74	0.64	0.54
$(BB^*)1^{++}$	0.92	0.80	0.66	0.58	0.50
$(B^*B^*)2^{++}$	0.94	0.82	0.67	0.59	0.51
$(B_sB_s)0^{++}$	—	—	1.42	1.01	0.74
$(B_sB_s^*)1^{++}$	—	1.77	1.03	0.81	0.63
$(B_s^*B_s^*)2^{++}$	—	1.81	1.05	0.83	0.64

The Belle collaboration has reported a $X(3930)$ state of 2^{++} at mass $M = 3929$ MeV[26], which is a candidate of the $c\bar{c}$ charmonium 2^3P_2 excited state χ'_{c2} . However its mass is about 40 MeV below the quark model calculations (in the Bhaduri model, the χ'_{c2} mass is $M = 3963.53$ MeV). Several authors have discussed the mass shifts of the coupled channel effect [27, 28, 29, 30]. If the above 2^{++} molecular state exists, the final state interaction will be important.

Now, we turn to the possible molecular states in other sectors. We observe that in $D_S^{(*)}D_S^{(*)}$ sector, there are no such molecular states due to the mass increase of m_s of the light quark pair. On the other hand, in the $B^{(*)}B^{(*)}$ sector, the binding energy increased as the mass increase of m_b of the heavy quark pair. So these bottomium-like molecular states with 1^{++} , 2^{++} , and even 0^{++} may also exist. We can also observe the $B_S^{(*)}B_S^{(*)}$ molecular states 1^{++} and 2^{++} if $f \sim -1.0$.

IV. SUMMARY

The quark model is extended by introducing the s -channel one gluon exchange interaction. The interaction has no effect on the inner quark structure of conventional $q\bar{q}$ mesons and qqq baryons. Since the interaction is short ranged — a smeared δ -interaction, the effect on the long ranged hadron-hadron interaction is expected to be very small. So the significant effect of this interaction is only in the so called hidden flavor states of multi-quark system.

We have calculated the heavy quark molecular states of $qQ\bar{Q}\bar{q}$ with $Q = c, b$ and $q = u, d, s$. We find that the interaction can be strong enough to bind the 1^{++} and 2^{++} states, and possibly 0^{++} states. To compare with the recent experiments, the $X(3872)$ is a candidate of the DD^* molecular state and the $X(3930)$ may be the χ'_{c2} state which couples to the D^*D^* 2^{++} molecular state. The calculation shows that it is more easy to bind the bottomium-like molecular states. Thus we expect that the similar bottomium-like molecular states of 1^{++} and 2^{++} should also exist.

Acknowledgments

We would like to thank Shi-Lin Zhu for useful discussions. This work was supported by the National Natural Science Foundation of China under Grants 10675008.

-
- [1] S. K. Choi et al. (Belle), Phys. Rev. Lett. **91**, 262001 (2003), hep-ex/0309032.
 - [2] D. E. Acosta et al. (CDF II), Phys. Rev. Lett. **93**, 072001 (2004), hep-ex/0312021.
 - [3] V. M. Abazov et al. (D0), Phys. Rev. Lett. **93**, 162002 (2004), hep-ex/0405004.
 - [4] B. Aubert et al. (BABAR), Phys. Rev. **D71**, 071103 (2005), hep-ex/0406022.
 - [5] F. E. Close and P. R. Page, Phys. Lett. **B578**, 119 (2004), hep-ph/0309253.
 - [6] M. B. Voloshin, Phys. Lett. **B579**, 316 (2004), hep-ph/0309307.
 - [7] E. S. Swanson, Phys. Lett. **B588**, 189 (2004), hep-ph/0311229.
 - [8] C.-Y. Wong, Phys. Rev. **C69**, 055202 (2004), hep-ph/0311088.
 - [9] N. A. Tornqvist, Phys. Lett. **B590**, 209 (2004), hep-ph/0402237.
 - [10] M. T. AlFiky, F. Gabbiani, and A. A. Petrov, Phys. Lett. **B640**, 238 (2006), hep-ph/0506141.
 - [11] S. Fleming, M. Kusunoki, T. Mehen, and U. van Kolck, Phys. Rev. **D76**, 034006 (2007), hep-ph/0703168.
 - [12] E. Braaten, M. Lu, and J. Lee, Phys. Rev. **D76**, 054010 (2007), hep-ph/0702128.
 - [13] C. Hanhart, Y. S. Kalashnikova, A. E. Kudryavtsev, and A. V. Nefediev, Phys. Rev. **D76**, 034007 (2007), 0704.0605.
 - [14] M. B. Voloshin, Phys. Rev. **D76**, 014007 (2007), 0704.3029.
 - [15] P. Colangelo, F. De Fazio, and S. Nicotri, Phys. Lett. **B650**, 166 (2007), hep-ph/0701052.
 - [16] M. Suzuki, Phys. Rev. **D72**, 114013 (2005), hep-ph/0508258.
 - [17] C. Meng, Y.-J. Gao, and K.-T. Chao (2005), hep-ph/0506222.
 - [18] S.-L. Zhu, Int. J. Mod. Phys. **E17**, 283 (2008), hep-ph/0703225.
 - [19] Y.-R. Liu, X. Liu, W.-Z. Deng, and S.-L. Zhu, Eur. Phys. J. **C56**, 63 (2008), 0801.3540.
 - [20] R. K. Bhaduri, L. E. Cohler, and Y. Nogami, Nuovo Cim. **A65**, 376 (1981).
 - [21] Z. Li, M. Guidry, T. Barnes, and E. S. Swanson (1994), hep-ph/9401326.
 - [22] H. Kameyama, M. Kamimura, and Y. Fukushima, Phys. Rev. **C40**, 974 (1989).
 - [23] K. Varga and Y. Suzuki, Phys. Rev. **A53**, 1907 (1996).

- [24] D. M. Brink and F. Stancu, Phys. Rev. **D57**, 6778 (1998).
- [25] C. Amsler et al. (Particle Data Group), Phys. Lett. **B667**, 1 (2008).
- [26] S. Uehara et al. (Belle), Phys. Rev. Lett. **96**, 082003 (2006), hep-ex/0512035.
- [27] Y. S. Kalashnikova, Phys. Rev. **D72**, 034010 (2005), hep-ph/0506270.
- [28] M. R. Pennington and D. J. Wilson, Phys. Rev. **D76**, 077502 (2007), 0704.3384.
- [29] B.-Q. Li, C. Meng, and K.-T. Chao, Phys. Rev. **D80**, 014012 (2009), 0904.4068.
- [30] P. G. Ortega, J. Segovia, D. R. Entem, and F. Fernandez (2009), 0907.3997.